

2014

MATHEMATICS

(Major)

Paper : 5.5

(Probability)

Full Marks : 60

Time : 3 hours

*The figures in the margin indicate full marks
for the questions*

1. Answer the following questions : 1×7=7
- (a) Write the sample space for the experiment of tossing a coin three times in succession or tossing three coins at a time.
 - (b) What conclusion one can make about the conditional probability $P(A|B)$ if $P(B) = 0$?
 - (c) Mention two properties which must be satisfied by the probability density function $f(x)$ of a continuous random variable X .

(2)

- (d) State the multiplication theorem of expectation.
- (e) State the relationship between the moment generating function of the sum of a number of independent random variables and the moment generating function of these individual random variables.
- (f) If X is a random variable and a, b are constants, then express variance of $(aX + b)$ in terms of variance of X .
- (g) Mention the relationship between the mean, median and mode of the normal distribution.

2. Answer the following questions : $2 \times 4 = 8$

- (a) Show that probability of any impossible event is zero.
- (b) If F is the distribution function of a random variable X and $a < b$, then show that

$$P(a < X \leq b) = F(b) - F(a)$$

- (c) If X and Y are two random variables, then show that

$$\text{cov}(X, Y) = E(XY) - E(X)E(Y)$$

(3)

- (d) If the random variable X follows binomial distribution with parameters n and p , then show that

$$E(X) = np$$

3. Answer any *three* parts from the following :

$5 \times 3 = 15$

- (a) If A and B are independent events, then show that \bar{A} and \bar{B} are also independent events.
- (b) A bag contains 6 white and 9 black balls. Four balls are drawn at a time. Find the probability for the first draw to give 4 white and the second to give 4 black balls if the balls drawn in the first draw are not replaced before the second draw.
- (c) A random variable X has the following probability function. Values of X —
- | | | | | | | | | |
|--------|-----|-----|------|------|------|-------|--------|------------|
| x | : 0 | 1 | 2 | 3 | 4 | 5 | 6 | 7 |
| $p(x)$ | : 0 | K | $2K$ | $2K$ | $3K$ | K^2 | $2K^2$ | $7K^2 + K$ |
- Find K and also evaluate $P(X < 6)$, $P(X \geq 6)$ and $P(0 < X < 5)$
- (d) A coin is tossed until a head appears. What is the expectation of the number of tosses required?

(4)

4. Answer any *three* parts from the following :

10×3=30

(a) (i) A probability curve $y = f(x)$ has a range from 0 to ∞ .

If $f(x) = e^{-x}$, find the mean and variance.

5

(ii) Let X be a continuous random variate with probability density function

$$\begin{aligned} f(x) &= ax, & 0 \leq x \leq 1 \\ &= a, & 1 \leq x \leq 2 \\ &= -ax + 3a, & 2 \leq x \leq 3 \\ &= 0, & \text{elsewhere} \end{aligned}$$

Compute $P(X \leq 1.5)$.

5

(b) (i) Show that expected value of X is equal to the expectation of the conditional expectation of X given Y i.e., $E(X) = E[E(X|Y)]$.

5

(ii) Discuss the effect of change of origin and scale on moment generating functions.

5

(c) (i) The mean and variance of a binomial distribution are 4 and $\frac{4}{3}$ respectively. Find $P(X \geq 1)$.

5

(ii) Ten coins are thrown simultaneously. Use binomial distribution to find the probability of getting at least seven heads.

5

(5)

(d) (i) Show that the mean and variance of the Poisson distribution are equal.

5

(ii) Write the probability density function of a random variable X which follows Normal distribution with mean μ and variance σ^2 . What is a standard normal variate? Find its mean and variance.

5
